



## International Journal of Molecular and Clinical Microbiology



### Short Communication

## Comparative of phenotypic tests in aerobic actinomycetes

Mehdi Fatahi-Bafghi

Department of Microbiology, Faculty of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.  
E-mail address: mehdifatahi@ssu.ac.ir

### Dear editor

I read with interest article that published entitled \*Isolation and identification of bioactive compound producing *Rhodococcus* spp. isolated from soil samples {IJMCM/5(1) (2015) 463-468}\* (Aghaei and Baserisalehi, 2015). Some of the genus such as *Nocardia*, *Gordonia*, *Mycobacterium* and *Rhodococcus* are in actinomycete family and they are Gram-positive and partially acid-fast. *Rhodococcus* species usually stain Gram-positive. Cells form as cocci or short rods which grow in length, and may form an extensively branched vegetative mycelium which may fragment. Microscopic aerial hyphae and spores are not usually produced. They are also non-motile. They are usually partially acid-fast due to the mycolic acid in their cell walls. Colonies of other rhodococci may be rough, smooth or mucoid and pigmented cream, buff, yellow, coral, orange or red. Although biochemical tests help to distinguish *Rhodococcus* from other organisms, differentiation from other aerobic actinomycetes can be difficult. Colonial and cell morphology cannot be used to distinguish among *Rhodococcus*, *Gordonia* and *Tsukamurella* species. I listed some of

phenotypic characterization of *Nocardia*, *Gordonia*, *Mycobacterium*, *Rhodococcus* and *Corynebacterium* in table 1 (Goodfellow, 1973; Goodfellow, 1974; Bell et al., 1998; Prescott 1991; Arenskötter et al., 2004; Stoecker et al., 2004; Li et al., 1994; Bafghi, 2015) and showed phenotypic tests such as microscopic examination, Gram and acid-fast staining, catalase, oxidase and motility tests that used by Aghaei et al is insufficient for the genus *Rhodococcus* confirmation. In literature, results of phenotypic tests are ambiguous for *Rhodococcus* identification and cannot distinguish the genus *Gordonia* of *Rhodococcus*. (Blanc et al., 2007) Authors do not explain about phenotypic tests results in this article and results are equivocal. Phenotypic tests with molecular methods such as sequence analysis of the 16S rRNA gene and PCR-RFLP (polymerase chain reaction-restriction fragment length polymorphism) should be better used for accurate identification in the genus level (Bell et al., 1998; Bell et al., 1999; Steingrube et al., 1997; Silva et al., 2012) that authors don't mentioned in article.

### Article history:

Received 23 December 2015

Accepted 3 July 2016

Available online 1 June 2016

**Table 1.** Various phenotypic tests in aerobic actinomycetes.

| Genus                               | soil | ISP5 agar | Bennet agar | Aerial hyphae | Gram | Acid-fast | Partially-acid fast | Motile | Catalase | Oxidase |
|-------------------------------------|------|-----------|-------------|---------------|------|-----------|---------------------|--------|----------|---------|
| <i>Nocardia</i>                     | +    | +         | +           | +             | +    | -*        | +                   | -      | +        | -       |
| <i>Gordonia</i>                     | +    | +         | +           | -             | +    | -*        | +                   | -      | +        | -       |
| <i>Mycobacterium</i> <sup>NTM</sup> | +/-  | -         | +/-         | -             | +    | +         | +                   | +/-    | +        | +       |
| <i>Rhodococcus</i>                  | +    | +         | +           | -             | +    | -*        | +                   | -      | +        | -       |
| <i>Corynebacterium</i>              | +/-  | +/-       | +/-         | -             | +    | -*        | +                   | +/-    | +/-      | +/-     |

\*In special circumstances are positive. NTM: non-tuberculosis mycobacterial

## Refereces

- Aghaei, S., Bahador, N., Baserisalehi, M., 2015. Isolation and identification of bioactive compound producing *Rhodococcus* spp. isolated from soil samples. International Journal of Molecular and Clinical Microbiology. 5(1): p. 463-468.
- Arenskötter, M., Bröker, D., and Steinbüchel, A., 2004. Biology of the metabolically diverse genus *Gordonia*. Applied and Environmental Microbiology. 70(6): p. 3195-3204.
- Bafghi, M.F., 2015. *Rhodococcus equi*: are phenotypic tests the best method for accurate identification? Comparative Clinical Pathology. 24(5): p. 979-979.
- Bell, K., et al., The genus *Rhodococcus*. Journal of Applied Microbiology, 1998. 85(2): p. 195-210.
- Bell, K., et al., 1999. Identification and environmental detection of *Rhodococcus* species by 16S rDNA targeted PCR. Journal of applied microbiology. 87(4): p. 472-480.
- Blanc, V., et al., 2007. *Gordonia terrae*: a difficult-to-diagnose emerging pathogen? Journal of clinical microbiology. 45(3): p. 1076-1077.
- Goodfellow, M., 1973. Characterisation of *Mycobacterium*, *Nocardia*, *Corynebacterium* and related taxa. Ann Soc Belg Med Trop. 53(4): p. 287-298.
- Goodfellow, M., et al., 1974. A co-operative numerical analysis of cultures considered to belong to the 'rhodochrous' taxon. Journal of general microbiology. 85(2): p. 291-302.
- Li, W.J., et al., 2004. *Nocardia alba* sp. nov., a novel actinomycete strain isolated from soil in China. Systematic and applied microbiology. 27(3): p. 308-312.
- Prescott, J.F., 1991. *Rhodococcus equi*: an animal and human pathogen. Clinical Microbiology Reviews. 4(1): p. 20-34.
- Silva, P.d., et al., 2012. Phenotypic and genotypic characterization of *Rhodococcus equi* isolated from sputum. Brazilian Journal of Infectious Diseases. 16(5): p. 409-415.
- Steingrube, V.A., et al., 1997. Rapid identification of clinically significant species and taxa of aerobic actinomycetes, including *Actinomadura*, *Gordonia*, *Nocardia*, *Rhodococcus*, *Streptomyces*, and *Tsukamurella* isolates, by DNA amplification and restriction endonuclease analysis. Journal of clinical microbiology. 35(4): p. 817-822.
- Stoecker, M.A., Herwig, R.P., and Staley, J.T., 1994. *Rhodococcus zopfii* sp. nov., a toxicant-degrading bacterium. International journal of systematic bacteriology. 44(1): p. 106-110.